



INDUSTRIAL  
MATHEMATICS  
INSTITUTE

2003:25

High order geometric smoothness  
for conservation laws

M. Campos-Pinto, A. Cohen and  
P. Petrushev

IMI

Preprint Series

Department of Mathematics  
University of South Carolina

# High order geometric smoothness for conservation laws <sup>\*</sup>

Martin Campos-Pinto, Albert Cohen, and Pencho Petrushev

## Abstract

The smoothness of the solutions of 1D scalar conservation laws is investigated and it is shown that if the initial value has smoothness of order  $\alpha$  in  $L^q$  with  $\alpha > 1$  and  $q = 1/\alpha$ , this smoothness is preserved at any time  $t > 0$  for the graph of the solution viewed as a function in a suitably rotated coordinate system. The precise notion of smoothness is expressed in terms of a scale of Besov spaces which also characterizes the functions that are approximated at rate  $N^{-\alpha}$  in the uniform norm by piecewise polynomials on  $N$  adaptive intervals. An important implication of this result is that a properly designed adaptive strategy should approximate the solution at the same rate  $N^{-\alpha}$  in the Hausdorff distance between the graphs.

## 1 Introduction

Solutions to hyperbolic equations derived from nonlinear conservation laws

$$\partial_t u + \text{Div}_x[f(u)] = 0, \quad u(x, 0) = u_0(x), \quad (1.1)$$

may develop discontinuities even if the initial data is smooth. This well known state of fact is the source of both theoretical difficulties - classical solutions should be replaced by weak solutions and side conditions need to be appended in order to ensure their uniqueness - as well as numerical difficulties

---

<sup>\*</sup>This work has been supported by the National Science Foundation Grant DMS-0200665 and by the European Community's Improving Human Potential Network HPRN-CT-202-00286 "Breaking Complexity".

- conventional discretization schemes may fail to converge and their convergence rate is in all cases limited by the lack of smoothness of the solution. We refer the reader to [10, 6, 7] for a general introduction to conservation laws.

In the case of scalar conservation laws, the classical theory developed by Kruzkov [8] ensures the uniqueness of an entropy solution  $u(x, t)$ . This solution is also stable in  $L^1$ , i.e.,

$$\|u(\cdot, t) - v(\cdot, t)\|_{L^1} \leq \|u_0 - v_0\|_{L^1} \quad (1.2)$$

for two solutions  $u$  and  $v$  with initial data  $u_0$  and  $v_0$ , and satisfies the BV diminishing property

$$\|u(\cdot, t)\|_{BV} \leq \|u_0\|_{BV}. \quad (1.3)$$

The BV boundedness plays a pivotal role in proving the convergence of numerical methods and deriving convergence rates with respect to the mesh size. As already mentioned, these rates are inherently limited by the lack of smoothness: the approximation  $u_h$  of a function  $u$  by piecewise polynomials on a uniform mesh cannot converge in  $L^1$  with a rate better than  $\mathcal{O}(h)$  when  $u$  has an isolated jump.

Adaptive methods offer a better compromise between error and number of degrees of freedom, especially when the solution is piecewise smooth with isolated singularities. From Approximation theory point of view these methods correspond to approximation from piecewise polynomials of a fixed degree on  $N$  intervals. Note that this is a nonlinear set since the  $N$  intervals may vary with the function being approximated, and therefore this type of approximation is referred to as *nonlinear approximation*. A precise description of those functions which can be approximated in  $L^1$  at rate  $N^{-\alpha}$  by such piecewise polynomial functions is given by the Besov space  $B_{q,q}^\alpha$  with  $1/q = 1 + \alpha$ , which consists of all functions  $u \in L^q$  such that

$$|u|_{B_{q,q}^\alpha}^q := \int_0^\infty [t^{-\alpha} \omega_k(u, t)_q]^q dt/t < \infty, \quad (1.4)$$

where  $k$  is an integer strictly larger than  $\alpha$  and  $\omega_k(u, t)_q := \sup_{|h| \leq t} \|\Delta_h^k u\|_{L^q}$  is the  $k$ -th order  $L^q$  modulus of smoothness. The norm in  $B_{q,q}^\alpha$  is defined by

$$\|u\|_{B_{q,q}^\alpha} := \|u\|_{L^q} + |u|_{B_{q,q}^\alpha}. \quad (1.5)$$

Roughly speaking, the functions in  $B_{q,q}^\alpha$  have  $\alpha$  derivatives in  $L^q$ . We refer to [2] as a general survey on nonlinear approximation.

In a series of papers [11, 3, 4], DeVore and Lucier have explored the smoothness properties of 1D scalar conservation laws using the above Besov spaces. They have shown that for all  $\alpha > 0$ , if the initial condition  $u_0$  belongs to  $B_{q,q}^\alpha$  with  $1/q = 1 + \alpha$ , then this property holds for the solution for all  $t > 0$ . The theorem of DeVore-Lucier shows that the solutions of conservation laws have an arbitrarily high order of smoothness  $\alpha > 0$  whenever the smoothness is measured in  $L^q$  with  $1/q = 1 + \alpha$ , and therefore  $q < 1$ . From a numerical perspective, it also indicates that a properly designed adaptive strategy should approximate the solution in  $L^1$  with an arbitrarily high rate of convergence with respect to the number of degrees of freedom. The proof of this theorem is based on the equivalence between smoothness and rate of nonlinear approximation, according to the following scheme:

1. The initial data  $u_0 \in B_{q,q}^\alpha$  is approximated at rate  $N^{-\alpha}$  by a piecewise polynomial function  $v_0$  on  $N$  intervals.
2. Then by the  $L^1$  stability (1.2) the solution  $u$  at time  $t > 0$  is approximated at the same rate  $N^{-\alpha}$  by the solution  $v$  with initial value  $v_0$ .
3. This rate of approximation allows to derive that  $u \in B_{q,q}^\alpha$ .

The main difficulty in this approach resides in the last step since it is no longer true that  $v$  is a piecewise polynomial on  $N$  intervals.

Since one of the goals of adaptive methods is to achieve uniformly accurate approximation, one could hope for similar results with the  $L^1$  norm replaced by the uniform ( $L^\infty$ ) norm as a measure of the error. However, such results are impossible since there is no stability in the uniform norm due to the development of discontinuities. A natural alternative is to measure the closeness between solutions and approximate solutions in the *Hausdorff distance between their completed graphs*, i.e.

$$d(u, v) = d_H(G_u, G_v),$$

where  $G_f$  denotes the completed graph of the function  $f$  and

$$d_H(A, B) := \max\left\{\sup_{a \in A} \inf_{b \in B} |a - b|, \sup_{b \in B} \inf_{a \in A} |a - b|\right\}$$

denotes the Hausdorff distance between the sets  $A$  and  $B$  (with  $|\cdot|$  denoting the Euclidean distance in  $\mathbb{R}^2$ ). Here the *completed graph*  $G_f$  of a function  $f$  is defined as the minimal closed set in  $\mathbb{R}^2$  which contains the graph of  $f$  and

is convex with respect to the  $y$ -direction, i.e., it is  $y$ -simple. It is easy to see that if  $f \in BV$  and  $f(x^-) \leq f(x) \leq f(x^+)$  for every  $x$ , then to obtain  $G_f$  one has to add to the graph of  $f$  every segment in the plane connecting the points  $(x, f(x^-))$  and  $(x, f(x^+))$  at every point  $x$ , where  $f$  is discontinuous (see [13]). The distance  $d(u, v)$  is a natural substitute for the  $L^\infty$  distance for discontinuous functions for two reasons: on the one hand it measures the closeness in  $L^\infty$  in regions where one of the functions is smooth enough since one easily checks that

$$\|u - v\|_{L^\infty} \leq d(u, v)[\|u'\|_{L^\infty} + 1],$$

and on the other hand it measures how accurately a sharp transition in  $u$  is matched in the  $x$ -direction by a sharp transition in  $v$ . In contrast to the  $L^\infty$  norm, stability results in the Hausdorff metric are available from [1], where it was recently proved that for 1D scalar conservation laws one has

$$d(u, v) \leq C(t)d(u_0, v_0) \tag{1.6}$$

with  $C(t) \sim 1 + t$ .

In this article, we shall use these results to establish high order smoothness results on the graph of the solution viewed as a function in a suitably rotated coordinate system. This approach is applicable in the case of strictly convex fluxes  $f$ , satisfying

$$0 < m \leq f''(u). \tag{1.7}$$

In a case like this, we invoke the Oleinik inequality which ensures that the entropy solution  $u$  of (1.1) satisfies at time  $t > 0$ ,

$$-\infty \leq u' \leq \frac{1}{mt}. \tag{1.8}$$

This inequality ensures that the graph of  $u$  is the graph of a Lipschitz function  $\tilde{u}$  in a suitably rotated coordinate system (which will be precisely specified in Section 3). In such a coordinate system the  $L^\infty$  distance between two solutions is equivalent to the Hausdorff distance between their graphs in the original coordinate system. This fact is illustrated on Figure 1.

We shall prove that the function  $\tilde{u}$  can be approximated in  $L^\infty$  by piecewise polynomials on  $N$  intervals at rate  $N^{-\alpha}$ , whenever  $u_0$  satisfies a similar property. As it will be explained in Section 2, the set of functions which can




Figure 1. Change of coordinate system.

be approximated in the uniform norm at rate (roughly)  $N^{-\alpha}$  with  $\alpha > 1$  by such piecewise polynomials is given by the space

$$\tilde{B}^\alpha := \{u \in W^{1,1}(\mathbb{R}) : u' \in B_{q,q}^{\alpha-1}, q = 1/\alpha\}, \quad (1.9)$$

The norm in  $\tilde{B}^\alpha$  is defined by

$$\|u\|_{\tilde{B}^\alpha} := \|u\|_{L^\infty} + \|u'\|_{B_{q,q}^{\alpha-1}}. \quad (1.10)$$

Notice that this space is slightly smaller than the Besov space  $B_{q,q}^\alpha$  which may contain discontinuous functions if  $q < 1$ .

We next state our main result.

**Theorem 1.1.** *Assume that  $u_0$  is a compactly supported function which satisfies  $u_0' \leq M$ . Then for all  $\alpha > 1$  and time  $t > 0$  the rotated solution  $\tilde{u}$  satisfies*

$$\|\tilde{u}\|_{\tilde{B}^\alpha} \lesssim \|u_0\|_{\tilde{B}^\alpha} + 1, \quad (1.11)$$

where the constant in  $\lesssim$  depends only on  $t$  and  $M$ .

From numerical perspective, this result indicates that a properly designed adaptive strategy should approximate the solution in the Hausdorff distance at an arbitrarily high rate with respect to the number of degrees of freedom.

The paper is organized as follows: In Section 2, we give some preliminary results for nonlinear approximation in  $L^\infty$  and on the Hausdorff stability of conservation laws. Using these results, we develop in Section 3 the strategy of DeVore-Lucier from [3, 4], namely, we construct approximate solutions which approximate the true solution at rate  $N^{-\alpha}$  in the Hausdorff metric, and as a consequence in  $L^\infty$  with respect to the rotated coordinate system. The “return ticket” which allows to derive the smoothness of  $\tilde{u}$  from the approximation rate relies on inverse estimates which are the objective of Section 4.

## 2 Preliminary results

### 2.1 Nonlinear piecewise polynomial approximation

For a fixed compact interval  $I$  and a positive integer  $k$ , let us denote by  $\Sigma_n$  the set of all piecewise polynomials of degree not exceeding  $k$  with no more than  $2^n$  pieces on  $I$ . Then for a given  $u \in L^p(I)$  ( $0 < p \leq \infty$ ) the error of best  $L^p$  approximation to  $u$  from  $\Sigma_n$  is defined by

$$\sigma_n(u)_p := \inf_{S \in \Sigma_n} \|u - S\|_{L^p}. \quad (2.1)$$

If some  $S_n$  realizes this infimum, it is said to be a best  $L^p$  approximation to  $u$  from  $\Sigma_n$ . We find useful the notion of a *near best* approximation, that corresponds to  $\|u - S_n\|_{L^p} \leq C\sigma_n(u)_p$  for some constant  $C \geq 1$  independent of  $n$  and  $u$ .

In order to describe the approximation rate, it is convenient to introduce the approximation space  $\mathcal{A}_q^\alpha(L^p)$ , defined as the set of all functions  $u \in L^p$  such that

$$\|u\|_{\mathcal{A}_q^\alpha(L^p)} := \left( \sum_{n=-1}^{\infty} [2^{n\alpha}\sigma_n(u)_p]^q \right)^{1/q} \quad (2.2)$$

is finite. Here we use the convention  $\Sigma_{-1} = \{0\}$ , so that  $\sigma_{-1}(u)_p := \|u\|_{L^p}$ . Clearly  $\mathcal{A}_\infty^\alpha(L^p)$  is the set of functions which are approximated in  $L^p$  by piecewise polynomials with accuracy  $\mathcal{O}(2^{-n\alpha})$ , and  $\mathcal{A}_q^\alpha(L^p)$  is a slight variation of this set since  $\mathcal{A}_\infty^{\alpha+\varepsilon}(L^p) \subset \mathcal{A}_q^\alpha(L^p) \subset \mathcal{A}_\infty^\alpha(L^p)$  for any  $\varepsilon > 0$ . We also recall that if  $\sigma_n(u)_p \rightarrow 0$  as  $n \rightarrow \infty$ , one obtains an equivalent norm in  $\mathcal{A}_q^\alpha(L^p)$  by replacing  $\sigma_n(u)_p$  by  $\|S_{n+1} - S_n\|_{L^p}$ , where  $S_n$  is a near-best approximation to  $u$  from  $\Sigma_n$ . Indeed, clearly  $\|S_{n+1} - S_n\|_{L^p} \lesssim \sigma_{n+1}(u)_p + \sigma_n(u)_p$  with a constant independent of  $n$ . On the other hand,  $S_n$  converges to  $u$  in  $L^p$  and hence  $\|u - S_n\|_{L^p}$  can be bounded by  $\sum_{n' \geq n} \|S_{n'+1} - S_{n'}\|_{L^p}$ , and we complete the argument by the discrete Hardy inequality.

Since the work of DeVore and Popov [5], it is known that when  $\alpha < k+1$ ,  $\mathcal{A}_q^\alpha(L^1)$  coincides with the Besov space  $B_{q,q}^\alpha$  with  $1/q = 1 + \alpha$ , and they have equivalent norms. In this article, we are interested in piecewise polynomial approximation of continuous functions in the uniform norm. In this context,  $\Sigma_n$  is redefined as the set of all *continuous* piecewise polynomials of degree  $\leq k$  with no more than  $2^n$  polynomial pieces. This type of approximation is studied by Petrushev in [12], where the following Jackson and Bernstein

estimates are established:

$$\sigma_n(u)_\infty \lesssim 2^{-\beta n} \|u'\|_{B_{r,r}^{\beta-1}}, \quad (2.3)$$

and

$$u \in \Sigma_n \Rightarrow \|u'\|_{B_{r,r}^{\beta-1}} \lesssim 2^{\beta n} \|u\|_{L^\infty}, \quad (2.4)$$

with  $1 < \beta < k + 1$  and  $r = 1/\beta$ . These estimates are the classical vehicle for characterizing the approximation spaces  $\mathcal{A}_q^\alpha(L^\infty)$  for  $0 < \alpha < \beta$  in terms of the real interpolation spaces  $(L^\infty, \tilde{B}^\beta)_{\frac{\alpha}{\beta}, q}$ , where

$$\tilde{B}^\beta := \{u : u' \in B_{r,r}^{\beta-1}, r = 1/\beta\}. \quad (2.5)$$

In the following, we shall prove directly that  $\mathcal{A}_q^\alpha(L^\infty)$  in fact coincides with  $\tilde{B}^\alpha$  for  $1 < \alpha < k + 1$ . As already mentioned  $\tilde{B}^\alpha$  is slightly smaller than  $B_{q,q}^\alpha$  and does not contain discontinuous functions.

**Lemma 2.1.** *We have  $\mathcal{A}_q^\alpha(L^\infty) = \tilde{B}^\alpha$ ,  $q = 1/\alpha$ , with equivalent norms.*

**Proof:** Assume that  $u \in \mathcal{A}_q^\alpha(L^\infty)$ , and denote by  $S_n$  ( $n \geq 0$ ) a near best  $L^\infty$  approximation to  $u$  from  $\Sigma_n$ . We consider the discontinuous piecewise polynomial  $T_n := S'_n$  of degree  $k - 1$  as an approximation to  $u'$ . Note that any polynomial  $S$  of degree  $k$  satisfies

$$\|S'\|_{L^1([a,b])} \leq C \|S\|_{L^\infty([a,b])},$$

where the constant  $C$  depends on  $k$ , but is independent of the interval  $[a, b]$  by a scaling argument. Since  $T_n - T_{n-1}$  is a piecewise polynomial on at most  $\frac{3}{2}2^n$  intervals  $I_j$ , we have

$$\|T_n - T_{n-1}\|_{L^1} \leq \sum_j \|T_n - T_{n-1}\|_{L^1(I_j)} \lesssim 2^n \|S_n - S_{n-1}\|_{L^\infty}.$$

This gives

$$\sum_{n=-1}^{\infty} [2^{n(\alpha-1)} \|T_n - T_{n-1}\|_{L^1}]^q \lesssim \|u\|_{\mathcal{A}_q^\alpha(L^\infty)}^q,$$

which in turn shows that  $T_n$  converges to an  $L^1$  function which is necessarily  $u'$ . It follows that

$$\|u'\|_{\mathcal{A}_q^{\alpha-1}(L^1)} \lesssim \|u\|_{\mathcal{A}_q^\alpha(L^\infty)},$$



and therefore, according to the result of [5] for piecewise polynomial approximation in  $L^1$ ,

$$\|u'\|_{B_{q,q}^{\alpha-1}(L^1)} \lesssim \|u\|_{\mathcal{A}_q^\alpha(L^\infty)}.$$

Now since  $\|u\|_{L^\infty} \leq \|u\|_{\mathcal{A}_q^\alpha(L^\infty)}$ , then

$$\|u\|_{\tilde{B}^\alpha} \lesssim \|u\|_{\mathcal{A}_q^\alpha(L^\infty)}.$$

For the estimate in the other direction, let us assume that  $u \in \tilde{B}^\alpha$ . Then  $u' \in B_{q,q}^{\alpha-1}$  with  $1/q = 1 + (\alpha - 1)$ , and due to the result of [5] for piecewise polynomial approximation in  $L^1$ , there exists a sequence  $(T_n)_{n \geq -1}$  of piecewise polynomials of degree  $k - 1$  with  $T_{-1} = 0$  such that  $T_n$  converges to  $u'$  in  $L^1$  and

$$\sum_{n=-1}^{\infty} 2^{(\alpha-1)qn} \|u' - T_n\|_{L^1}^q \lesssim \|u'\|_{B_{q,q}^{\alpha-1}}^q.$$

Clearly, there is a subdivision with at most  $2^{n+1}$  intervals  $I_j$  such that  $T_n$  is a polynomial on each of them and

$$\|u' - T_n\|_{L^1(I_j)} \leq 2^{-n} \|u' - T_n\|_{L^1}.$$

On each interval  $I_j = [a_j, b_j]$ , we define

$$P_{n+1}(x) := u(a_j) + \int_{a_j}^x T_n(s) ds, \quad (2.6)$$

and further modify  $P_{n+1}$  into

$$S_{n+1}(x) := P_{n+1}(x) + (u(b_j) - P_{n+1}(b_j)) \frac{x - a_j}{b_j - a_j}. \quad (2.7)$$

Thus the resulting  $S_{n+1}$  is in  $\Sigma_{n+1}$ . On each  $I_j$ , we clearly have

$$|u(x) - P_{n+1}(x)| \leq \|u' - T_n\|_{L^1(I_j)} \leq 2^{-n} \|u' - T_n\|_{L^1},$$

and hence

$$|u(b_j) - P_{n+1}(b_j)| \frac{x - a_j}{b_j - a_j} \leq 2^{-n} \|u' - T_n\|_{L^1}.$$

Consequently,

$$\|u - S_{n+1}\|_{L^\infty} \leq 2^{-n+1} \|u' - T_n\|_{L^1},$$

which implies

$$\|u\|_{\mathcal{A}_q^\alpha(L^\infty)}^q \lesssim \|u\|_{L^\infty}^q + \|u'\|_{\mathcal{A}_q^{\alpha-1}(L^1)}^q.$$

Now invoking the result of [5] for piecewise polynomial approximation in  $L^1$ , we conclude

$$\|u\|_{\mathcal{A}_q^\alpha(L^\infty)} \lesssim \|u\|_{\tilde{B}^\alpha}.$$

The proof is complete.  $\square$

In the second part of the proof of Lemma 2.1, we constructed the approximation  $S_{n+1}$  to  $u$  by using that  $T_n$  approximates  $u'$  (see (2.6)-(2.7)). For future use, it will be useful to construct  $S_n$  so that if  $u' \leq M$ , then  $S_n$  also satisfies  $S'_n \leq M$ . To this end, we slightly modify the above construction as is described in the following. Once the intervals  $I_j$  are determined, we define on each of them a new approximation  $R_{n+1}$  to  $u'$  as the orthogonal projection of  $u'$  onto the polynomials of degree  $k-1$ , namely,  $R_{n+1}$  is defined on each  $I_j$  so that

$$\int_{I_j} [R_{n+1}(x) - u'(x)] x^\nu dx = 0, \quad \nu = 0, \dots, k-1.$$

Since this orthogonal projection is a near best  $L^1$  approximation, we have

$$\|u' - R_{n+1}\|_{L^1(I_j)} \lesssim \|u' - T_n\|_{L^1(I_j)}.$$

Inside  $I_j$ , there are at most  $[k/2+1]$  disjoint intervals on which  $R_{n+1}(x) > M$ . On each of them we replace  $R_{n+1}$  by the constant  $M$  and on the remaining part  $\tilde{I}_j$  of  $I_j$  we modify  $R_{n+1}$  as  $M - c(M - R_{n+1})$ , where  $c$  ensures that the integral of  $R_{n+1}$  on  $I_j$  remains unchanged. Note that since this integral is

$$\int_{I_j} R_{n+1} = \int_{I_j} u' \leq M|I_j|,$$

then the constant

$$c := \frac{\int_{I_j} [M - R_{n+1}]}{\int_{\tilde{I}_j} [M - R_{n+1}]}$$

is necessarily in  $[0, 1]$ , and consequently  $M - c(M - R_{n+1}) \leq M$  on  $\tilde{I}_j$ . The resulting function  $U_{n+a}$  has at most  $2^{n+a}$  pieces with  $a = 1 + [\log_2 k]$  and

satisfies  $U_{n+a} \leq M$  everywhere. We finally remark that this modification can only improve the  $L^1$  approximation error on  $I_j$ . Indeed, on the one hand

$$\|u' - U_{n+a}\|_{L^1(I_j \setminus \tilde{I}_j)} \leq \|u' - R_{n+1}\|_{L^1(I_j \setminus \tilde{I}_j)} - \int_{I_j \setminus \tilde{I}_j} [R_{n+1} - M],$$

and on the other hand

$$\begin{aligned} \|u' - U_{n+a}\|_{L^1(\tilde{I}_j)} &= \|u' - M - c(R_{n+1} - M)\|_{L^1(\tilde{I}_j)} \\ &\leq \|u' - R_{n+1}\|_{L^1(\tilde{I}_j)} + (1 - c)\|M - R_{n+1}\|_{L^1(\tilde{I}_j)} \\ &= \|u' - R_{n+1}\|_{L^1(\tilde{I}_j)} + (\int_{\tilde{I}_j} [M - R_{n+1}] - \int_{I_j} [M - R_{n+1}]) \\ &= \|u' - R_{n+1}\|_{L^1(\tilde{I}_j)} + \int_{I_j \setminus \tilde{I}_j} [R_{n+1} - M]. \end{aligned}$$

Therefore

$$\|u' - U_{n+a}\|_{L^1(I_j)} \leq \|u' - R_{n+1}\|_{L^1(I_j)} \lesssim \|u' - T_n\|_{L^1(I_j)}. \quad (2.8)$$

We now define  $S_{n+a} \in \Sigma_{n+a}$  on each interval  $I_j$  by

$$S_{n+a}(x) := u(a_j) + \int_{a_j}^x U_{n+a}(s) ds. \quad (2.9)$$

The continuity of  $S_{n+a}$  is ensured since by construction  $\int_{I_j} U_{n+a} = \int_{I_j} u'$  and we clearly have  $S'_{n+a} \leq M$ . We complete the argument as in the proof of Lemma 2.1, namely, we have

$$\|u - S_{n+a}\|_{L^\infty} \lesssim 2^{-n} \|u' - T_n\|_{L^1},$$

and hence

$$\|u\|_{\mathcal{A}_q^\alpha(L^\infty)} \leq \left( \sum_{n=-1}^{\infty} [2^{n\alpha} \|u - S_n\|_{L^\infty}]^q \right)^{1/q} \lesssim \|u\|_{\tilde{B}^\alpha}, \quad (2.10)$$

where  $S_n := 0$  for  $-1 \leq n < a$ .

## 2.2 Hausdorff stability and rotated graphs

In [1], it is proved that scalar conservation laws are stable in the Hausdorff metric  $d(\cdot, \cdot)$  with respect to perturbations of the initial condition. More

precisely, if  $u$  and  $v$  are solutions of (1.1) with initial values  $u_0$  and  $v_0$ , and if for some  $M > 0$  the initial condition  $u_0$  satisfies

$$u'_0 \leq M \text{ or } u'_0 \geq -M, \quad (2.11)$$

then we have

$$d(u, v) \leq C(t)d(u_0, v_0), \quad t > 0, \quad (2.12)$$

with  $C(t) \sim 1 + M(1 + t)$ . A stability result is also established with respect to a perturbation of the flux function: If  $u$  and  $v$  are solutions of (1.1) with initial value  $u_0$  and fluxes  $f$  and  $g$ , respectively, then at time  $t > 0$ , we have

$$d(u, v) \leq C(t)\|f' - g'\|_{L^\infty} \quad (2.13)$$

with  $C(t) \sim 1 + t$ . These two results can be combined, namely, if  $u$  and  $v$  are solutions of (1.1) with initial value  $u_0$  and  $v_0$  and fluxes  $f$  and  $g$ , and if  $u_0$  satisfies (2.11), then

$$d(u, v) \leq C(t)[d(u_0, v_0) + \|f' - g'\|_{L^\infty}] \quad (2.14)$$

with  $C(t) \sim 1 + M(1 + t)$ .

As already explained in the introduction, our main idea is by employing the Oleinik inequality (1.8) to replace the Hausdorff distance by the  $L^\infty$  distance in a suitably rotated coordinate system. Indeed, assuming that  $u$  satisfies (1.8), it is readily seen that the graph of  $u$  is also the graph of a Lipschitz function  $\bar{u}$  in the rotated coordinate system defined by

$$\begin{cases} \bar{x} = cx - sy \\ \bar{y} = sx + cy \end{cases} \quad (2.15)$$

with  $c := \cos \theta$ ,  $s := \sin \theta$ ,  $\theta \in [0, \pi/2[$  such that

$$\tau := s/c = \tan \theta = mt/2. \quad (2.16)$$

One can indeed readily check that

$$-\tau^{-1} \leq \bar{u}'(\bar{x}) \leq 2\tau + \tau^{-1}. \quad (2.17)$$

Clearly, the rotated solution  $\bar{u}$  is not compactly supported since it coincides with the function  $\bar{y} = \tau\bar{x}$  outside the region corresponding to the support

of  $u$ . In order to preserve the compactness of the support, we modify  $\bar{u}$  by setting

$$\tilde{u} := \bar{u} - \tau \bar{x}. \quad (2.18)$$

Thus the new coordinate system is

$$\begin{cases} \tilde{x} = \bar{x} = cx - sy \\ \tilde{y} = c^{-1}y. \end{cases} \quad (2.19)$$

If  $u$  is supported on  $I(t) = [a(t), b(t)]$ , then  $\tilde{u}$  is supported on  $\tilde{I}(t) = [ca(t), cb(t)]$ . Clearly, we still have a Lipschitz bound

$$|\tilde{u}(\tilde{x}) - \tilde{u}(\tilde{y})| \leq \nu |\tilde{x} - \tilde{y}| \quad (2.20)$$

with

$$\nu := \tau + \tau^{-1}. \quad (2.21)$$

We also remark that if  $u \in BV$ , then  $\tilde{u} \in BV$ , and

$$|\tilde{u}|_{BV(\tilde{I})} \leq c^{-1} |u|_{BV(I)}, \quad (2.22)$$

which follows immediately from the definition of the total variation:

$$|u|_{BV} := \sup \sum_{i=1}^n |u(x_i) - u(x_{i-1})|,$$

where the supremum is taken over all selections of points  $x_0 < \dots < x_n$  in the support of  $u$ .

It is easy to see that if  $\tilde{u}$  and  $\tilde{v}$  are obtained from  $u$  and  $v$  by such a change of the coordinate system, then

$$\|\tilde{u} - \tilde{v}\|_{L^\infty} = \|\bar{u} - \bar{v}\|_{L^\infty} \leq (1 + \nu)d(\bar{u}, \bar{v}) = (1 + \nu)d(u, v),$$

and in the other direction,

$$d(u, v) = d(\bar{u}, \bar{v}) \leq \|\bar{u} - \bar{v}\|_{L^\infty} = \|\tilde{u} - \tilde{v}\|_{L^\infty}.$$

Therefore, the Hausdorff distance between two solutions is equivalent to the  $L^\infty$  distance between the rotated solutions. In particular, if  $u$  and  $v$  are solutions of (1.1) with initial values  $u_0$  and  $v_0$  and fluxes  $f$  and  $g$ , and if  $u_0$  satisfies (2.11), then we have

$$\|\tilde{u} - \tilde{v}\|_{L^\infty} \leq C(t)[\|u_0 - v_0\|_{L^\infty} + \|f' - g'\|_{L^\infty}] \quad (2.23)$$

with  $C(t) \sim \nu[1 + M(1 + t)]$ .

### 3 Proof of the regularity theorem

The proof of Theorem 1.1 relies on an approximation procedure by piecewise algebraic functions which stay close to the solution  $u$  in the Hausdorff metric for all  $t > 0$ . As shown above, this stability will hold in  $L^\infty$  in the coordinate system (2.19).

#### 3.1 Approximate solutions

Assuming that  $u_0 \in \tilde{B}^\alpha$  satisfies  $u'_0 \leq M$ , let  $S_n$  be the  $L^\infty$  approximation to  $u_0$  defined in (2.9). We recall that  $S_n$  is made up of at most  $2^n$  polynomial pieces of degree  $\leq k$  with  $k > \alpha - 1$ , and that it satisfies

$$S'_n \leq M. \quad (3.1)$$

We also observe that since  $S'_n = U_n$  is a near best  $L^1$  approximation of  $u'_0$ , then

$$\|S'_n\|_{L^1} \leq C\|u'_0\|_{L^1}$$

for some constant  $C$ , and therefore

$$\|S_n\|_{BV} \leq C\|u_0\|_{BV}. \quad (3.2)$$

Notice that  $S_n$  is not necessarily a near best  $L^\infty$  approximation to  $u_0$ . However, (2.10) guarantees that it is good enough for our purposes. Clearly, there is an interval  $\Omega$  (whose size may depend on  $\|u_0\|_{BV}$ ) such that  $u_0(x)$  and  $S_n(x)$  belong to  $\Omega$  for any  $x$ .

We next approximate the flux function. Assume that  $f \in C^2$  and  $f$  is strictly convex, so that there exist two constants  $m$  and  $\bar{m}$  such that

$$0 < m \leq f'' \leq \bar{m} \quad \text{on } \Omega.$$

We also assume that  $f$  belongs to  $W^{r+1,\infty}(\Omega)$ . Then by a classical spline approximation result, there exists an  $r - 1$  times continuously differentiable piecewise polynomial function  $g_n$  of degree  $\leq r$  with uniform knots at the points  $j2^{-n}$ ,  $j \in \mathbb{Z}$ , such that

$$\|f^{(l)} - g_n^{(l)}\|_{L^\infty(\Omega)} \leq C2^{-n(r+1-l)}\|f^{(r+1)}\|_{L^\infty(\Omega)} \quad \text{for } l = 0, \dots, r. \quad (3.3)$$

Changing slightly the constants  $m$  and  $\bar{m}$ , we may assume that the functions  $g_n$  also satisfy

$$0 < m \leq g_n'' \leq \bar{m} \quad \text{on } \Omega. \quad (3.4)$$

We now define  $s_n$  as the entropy solution at time  $t$  of (1.1) with initial value  $S_n$  and flux  $g_n$ , and denote it by  $\tilde{s}_n$  in the coordinate system (2.19). Before going any further, we observe that our stability result (2.23) combined with (3.3) guarantees that

$$\|\tilde{u} - \tilde{s}_n\|_{L^\infty} \leq C(t)[\|u_0 - S_n\|_{L^\infty} + 2^{-nr}] \quad (3.5)$$

as well as

$$\|\tilde{s}_{n+1} - \tilde{s}_n\|_{L^\infty} \leq C(t)[\|S_{n+1} - S_n\|_{L^\infty} + 2^{-nr}]. \quad (3.6)$$

Therefore,  $\tilde{s}_n$  approximates  $\tilde{u}$  with the same rate as  $S_n$  approximates  $u_0$ , up to an additional term  $2^{-nr}$ . In the following, we assume that  $\alpha + 1 < r$ . In particular, we can set  $r := k + 2$ .

## 3.2 Structure of the approximate solutions

We recall that a function  $y := y(x)$  is said to be *algebraic* on an interval  $J$  if there exists a polynomial  $F$  in two variables such that  $F(x, y(x)) = 0$  for  $x \in J$ . We shall now describe the structure of the approximate solutions  $\tilde{s}_n$  in terms of particular algebraic pieces  $(y, J)$ .

**Lemma 3.1.** *There exists a partition of the support of  $\tilde{s}_n$  into  $\mathcal{O}(2^n)$  intervals such that on each interval  $J$ , the function  $\tilde{s}_n$  coincides with an algebraic piece  $(y, J)$  of one of the following two types:*

**Type I:**  $y$  satisfies  $\|y'\|_{L^\infty(J)} \leq \nu$  and the algebraic equation

$$R(T(x)) = y(x) + \nu x, \quad x \in J, \quad (3.7)$$

where  $\nu$  is defined in (2.21),  $T(x) := y(x) + \nu x - Q(y(x))$ , and  $R$  and  $Q$  are algebraic polynomials of degrees  $k(r - 1)$  and  $r - 1$ , satisfying

$$\begin{aligned} (\mathbf{A}_1) \quad & 2 \leq Q' \leq c_1 \quad \text{on } y(J), \\ (\mathbf{A}_2) \quad & 0 < R' \leq c_2 \quad \text{on } T(J), \end{aligned}$$

for two constants  $c_1$  and  $c_2$ .

**Type II:**  $y$  satisfies

$$y(0) = y(x) + \nu x, \quad x \in J, \quad (3.8)$$

i.e.,  $\tilde{s}_n$  is affine on  $J$  with slope  $-\nu$ .

**Proof:** Following DeVore-Lucier [4], we begin by introducing two special types of points. First, let  $\{a_i\}_{0 \leq i \leq A}$  denote the knots of  $S_n$ , that is, the points where  $S_n$  changes from one polynomial piece to another. By construction,  $A \leq 2^n$ . Let then  $\{b_i\}_{0 \leq i \leq B}$  denote the *isolated* points such that  $S_n(b_i)$  is a knot of  $g_n$ , that is,  $S_n(b_i) = j2^{-n}$  for some  $j$ . To count them, we shall denote by  $\{\tilde{b}_i\}_{0 \leq i \leq \tilde{B}}$  all  $b_j$ 's such that  $S_n(b_{j-1}) = S_n(b_j)$ , and we denote the remaining ones by  $\{\bar{b}_i\}_{0 \leq i \leq \bar{B}}$ . Now, we have  $\text{Var}_{[\bar{b}_i, \bar{b}_{i+1}]}(S_n) \geq 2^{-n}$  for each  $i$ , hence  $\|S_n\|_{BV} \geq \sum_{i=0}^{\bar{B}-1} \text{Var}_{[\bar{b}_i, \bar{b}_{i+1}]}(S_n) \geq \bar{B} 2^{-n}$ , and we infer from (3.2) that  $\bar{B} \lesssim \|u_0\|_{BV} 2^n$ . On the other hand, if  $I_j$  is an interval where  $S_n$  coincides with the polynomial  $P_j$ ,  $P_j'$  should vanish at least once in each  $[\tilde{b}_i, \tilde{b}_{i+1}] \subset I_j$ . Since  $P_j'$  is of degree not exceeding  $k$  and by definition there are no second type points in  $I_j$  when  $P_j$  is a constant, we see that  $\tilde{B}$  is of order  $\mathcal{O}(2^n)$ , and so is  $B$ .

In [9], Lax shows that if the initial data  $S_n$  is continuous and the flux function  $g_n$  is strictly convex, the entropy solution  $s_n$  of (1.1) satisfies

$$s_n(x, t) = S_n(z), \text{ where } z := z(x, t) \text{ is a solution of } \frac{x - z}{t} = g_n'(S_n(z)).$$

There may be many solutions of this equation, but a minimization property picks a specific value  $z(x, t)$ . Lax shows that  $z(x, t)$  is an increasing function of  $x$  for a fixed  $t$ . Shocks occur wherever  $z(x, t)$  is discontinuous in  $x$ . If we denote by  $\sigma_i$  the positions of these shocks and set  $z_i^- := z(\sigma_i^-, t)$  and  $z_i^+ := z(\sigma_i^+, t)$ , this means that the function

$$\mathcal{S} : z \rightarrow z + t g_n'(S_n(z)) \tag{3.9}$$

is increasing on each interval  $[z_i^+, z_{i+1}^-]$ , while  $\mathcal{S}(z_i^-) = \mathcal{S}(z_i^+) = \sigma_i$ . From our previous discussion, we can describe  $\mathcal{S}$  as  $\mathcal{O}(2^n)$  polynomial pieces of degree at most  $k(r-2)$ , so it follows that there cannot be more than  $\mathcal{O}(2^n)$  shocks. In addition, we see that there is a partition  $\{I_i^0\}_{1 \leq i \leq C 2^n}$  such that  $\mathcal{S}$  is an increasing polynomial on each interval  $I_i^0$  and satisfies

$$s_n(\mathcal{S}(z)) = S_n(z), \quad z \in I_i^0 \tag{3.10}$$

(here  $s_n$  is multivalued at the shocks), while the intervals  $I_i^t := \mathcal{S}(I_i^0)$  recover  $\mathbb{R}$  and overlap only at the boundaries. Writing  $x = \mathcal{S}(z)$ , this leads to

$$\mathcal{S}(x - t g_n'(s_n(x))) = x, \quad x \in I_i^t. \tag{3.11}$$



Finally, we observe that in the coordinate system (2.19) each algebraic piece  $(s_n, I_i^t)$  becomes a piece of Type I, while the shocks become pieces of Type II, as is seen from Figure 1. Indeed, let us fix  $i$  and let  $P$  and  $Q$  denote the polynomials coinciding with  $c^{-1}S_n(s \cdot)$  and  $s^{-1}tg_n'(c \cdot)$  on  $s^{-1}I_i^0$  and  $c^{-1}S_n(I_i^0)$  respectively. Define also  $R := Id + Q \circ P$  the polynomial which coincides with  $s^{-1}\mathcal{S}(s \cdot)$  on  $s^{-1}I_i^0$ . After a little algebra, in the new coordinate system (3.11) becomes

$$R(\tilde{s}_n(\tilde{x}) + \nu\tilde{x} - Q(\tilde{s}_n(\tilde{x}))) = \tilde{s}_n(\tilde{x}) + \nu\tilde{x},$$

which gives (3.7) with  $J := \tilde{I}_i^t$ . Then

$$Q' = \frac{t}{\tau} g_n''(c \cdot) \quad \text{and} \quad R' = 1 + t g_n''(S_n(s \cdot)) S_n'(s \cdot),$$

and hence **(A<sub>1</sub>)**–**(A<sub>2</sub>)** follow readily from (3.4) and (3.1) with  $c_1 = 2\tilde{m}/m$  and  $c_2 = 1 + t\tilde{m}M$ .  $\square$

### 3.3 An inverse estimate

According to Lemma 3.1, each difference  $\tilde{s}_n - \tilde{s}_{n-1}$  is made of  $\mathcal{O}(2^n)$  algebraic pieces  $(A, J)$  which are differences of pieces of first or second type. Following DeVore and Lucier (Lemma 4.2 in [4]), we can further split these pieces in order to obtain a partition consisting of  $\mathcal{O}(2^n)$  pieces  $(A, J)$ , each of them monotone together with all its derivatives of order  $\leq k + 1$ . We next state an inverse estimate for such pieces which will allow to complete the proof of Theorem 1.1.

**Lemma 3.2.** *If  $(A, J)$  is an algebraic piece of  $\tilde{s}_n - \tilde{s}_{n-1}$ , then*

$$\|A' \cdot \mathbb{1}_J\|_{B_{q,q}^{\alpha-1}} \lesssim \|A\|_{L^\infty(J)} + 2^{-(r-1)n} \quad (3.12)$$

with a constant independent of  $n$ .

This inverse estimate has a delicate proof which will be given in Section 4.

From (3.12), we next deduce an inverse inequality for the functions  $\tilde{s}_n - \tilde{s}_{n-1}$ . Assuming that  $\{(A_i, J_i)\}_{1 \leq i \leq C 2^n}$  is a subdivision of  $\tilde{s}_n - \tilde{s}_{n-1}$  into algebraic pieces, we observe that the continuity of each  $\tilde{s}_n$  yields

$$\tilde{s}'_n - \tilde{s}'_{n-1} = \sum_{i=1}^{C 2^n} A'_i \cdot \mathbb{1}_{J_i}.$$

Therefore, using the  $q$ -triangle inequality for  $B_{q,q}^{\alpha-1}$ , we have

$$\begin{aligned} \|\tilde{s}'_n - \tilde{s}'_{n-1}\|_{B_{q,q}^{\alpha-1}}^q &\leq \sum_{i=1}^{C2^n} \|A'_i \mathbb{1}_{J_i}\|_{B_{q,q}^{\alpha-1}}^q \\ &\lesssim \sum_{i=1}^{C2^n} [\|A_i\|_{L^\infty(J_i)} + 2^{-n(r-1)}]^q \\ &\lesssim 2^n \|\tilde{s}_n - \tilde{s}_{n-1}\|_{L^\infty}^q + 2^{-n((r-1)q-1)} \end{aligned} \quad (3.13)$$

and, using (3.6), it follows that

$$\|\tilde{s}'_n - \tilde{s}'_{n-1}\|_{B_{q,q}^{\alpha-1}}^q \lesssim 2^n \|S_{n+1} - S_n\|_{L^\infty}^q + 2^{-n((r-1)q-1)}.$$

From (3.5), it also appears that  $\tilde{u}$  can be decomposed into a telescopic sum

$$\tilde{u} = \sum_{n=0}^{\infty} \tilde{s}_n - \tilde{s}_{n-1}.$$

Then applying again the  $q$ -triangle inequality, we obtain

$$\begin{aligned} \|\tilde{u}'\|_{B_{q,q}^{\alpha-1}}^q &\leq \sum_{n=0}^{\infty} \|\tilde{s}'_n - \tilde{s}'_{n-1}\|_{B_{q,q}^{\alpha-1}}^q \\ &\lesssim \sum_{n=0}^{\infty} [2^n \|S_n - S_{n-1}\|_{L^\infty}^q + 2^{-n((r-1)q-1)}] \\ &\lesssim \|u_0\|_{\tilde{B}^\alpha}^q + 1, \end{aligned}$$

where we used our assumption  $r - 1 > \alpha = 1/q$ . The proof of Theorem 1.1 is thus complete except for the proof of Lemma 3.2.

## 4 Proof of the inverse estimate

In this section,  $n$  is a fixed positive integer and  $(A, J)$  denotes an algebraic piece of  $\tilde{s}_n - \tilde{s}_{n-1}$ .

### 4.1 An intermediate estimate

In order to prove Lemma 3.2, we first establish the following intermediate inverse inequality.

**Lemma 4.1.** *If  $(A, J)$  is an algebraic piece of  $\tilde{s}_n - \tilde{s}_{n-1}$ , then*

$$\|A'\|_{L^\infty(J)} \lesssim |J|^{-1} \left( \|A\|_{L^\infty(J)} + 2^{-(r-1)n} \right) \quad (4.1)$$

*with a constant independent of  $n$ .*

**Proof:** We let  $y(x)$  and  $\bar{y}(x)$  denote the algebraic pieces of  $\tilde{s}_n$  and  $\tilde{s}_{n-1}$  on the interval  $J$ . Several cases are possible, depending on whether  $y$  and  $\bar{y}$  are of Type I or Type II. However, we observe that there is nothing to prove when  $y$  and  $\bar{y}$  are both of Type II. Thus we can always assume that  $y$  is of Type I and set

$$\Theta(x) := 1 - R'(T)(1 - Q'(y)).$$

We begin by establishing the equivalences

$$|\Theta(x)| \sim 1, \quad x \in J, \quad (4.2)$$

and

$$|T(J)| \sim |J| \quad (4.3)$$

with constants of equivalence independent of  $n$ .

For the proof of (4.2), we first see using  $(\mathbf{A}_1)$ – $(\mathbf{A}_2)$  that  $\|\Theta\|_{L^\infty(J)} \leq 1 + c_2(1 + c_1)$ . In the other direction, differentiating both sides of (3.7) and the expression for  $T(x)$  with respect to  $x$  yields

$$R'(T)T'(x) = y'(x) + \nu \quad (4.4)$$

and

$$T'(x) = \nu - y'(x)[Q'(y) - 1]. \quad (4.5)$$

Hence

$$y'(x)\Theta(x) = \nu[R'(T) - 1]. \quad (4.6)$$

Let  $J_+ := \{x \in J; |1 - R'(T)| \geq 1/2\}$  and  $J_- := J \setminus J_+$ . If  $x \in J_+$ , then  $|y'(x)\Theta(x)| \geq \nu/2$ , and using  $|y'(x)| \leq \nu$  it follows that  $|\Theta(x)| \geq 1/2$ . In the case when  $x \in J_-$ , we infer from the positivity of  $R'(T)$  on  $J$  that  $1/2 > 1 - R'(T)$ , and using  $(\mathbf{A}_1)$ , it follows that

$$\begin{aligned} |\Theta(x)| &\geq R'(T)Q'(y) - |1 - R'(T)| \\ &\geq (1/2)Q'(y) - 1/2 \\ &\geq 1/2. \end{aligned}$$

Hence  $|\Theta(x)| \geq 1/2$  for  $x \in J$  and the proof of (4.2) is complete.

We turn to the proof of (4.3). From (4.5), it is clear that  $\|T'\|_{L^\infty(J)} \leq \nu(2 + c_1)$ . To bound  $T'(x)$  from below, suppose first that  $y'(x) \geq 0$ . Then (4.4) together with  $(\mathbf{A}_2)$  yields  $T'(x) \geq \nu/c_2$ . If  $y'(x) \leq 0$ , then (4.5) along with  $(\mathbf{A}_1)$  implies  $T'(x) \geq \nu$ , and (4.3) follows.

We recall the following classical inequalities, valid for arbitrary intervals  $G, G'$  such that  $G \subset G'$ , and a polynomial  $P$  of degree  $\leq l$ :

$$(\mathbf{P}_1) \quad \|P\|_{L^\infty(G')} \leq C \left( \frac{|G'|}{|G|} \right)^l \|P\|_{L^\infty(G)},$$

$$(\mathbf{P}_2) \quad \|P'\|_{L^\infty(G)} \leq C |G|^{-1} \|P\|_{L^\infty(G)}.$$

We now consider the case where  $\bar{y}$  is of Type II. By (4.6), (4.2), and (3.8), we have

$$\begin{aligned} \|y' - \bar{y}'\|_{L^\infty(J)} &= \nu \|\Theta^{-1}(R'(T) - 1) + 1\|_{L^\infty(J)} \\ &= \nu \|\Theta^{-1}R'(T)Q'(y)\|_{L^\infty(J)} \\ &\lesssim \|R'(T)\|_{L^\infty(J)} \\ &\lesssim \|R'\|_{L^\infty(T(J))} \\ &\lesssim |T(J)|^{-1} \|R - \bar{y}(0)\|_{L^\infty(T(J))} \\ &\lesssim |J|^{-1} \|R(T) - \bar{y}(0)\|_{L^\infty(J)} \\ &\lesssim |J|^{-1} \|y - \bar{y}\|_{L^\infty(J)}, \end{aligned}$$

which proves the lemma in this case. Here the first inequality is again (4.2) together with  $(\mathbf{A}_1)$ , the third one is  $(\mathbf{P}_2)$ , the fourth one is (4.3), and the last one is (3.7) together with (3.8).

Let now  $y$  and  $\bar{y}$  be both of Type I. We use (4.6), (4.2), and  $(\mathbf{A}_2)$  to obtain

$$\begin{aligned} \|y' - \bar{y}'\|_{L^\infty(J)} &= \nu \|(\Theta \bar{\Theta})^{-1}[\bar{\Theta}(R'(T) - 1) - \Theta(\bar{R}'(\bar{T}) - 1)]\|_{L^\infty(J)} \\ &\lesssim \|\bar{\Theta}(R'(T) - 1) - \Theta(\bar{R}'(\bar{T}) - 1)\|_{L^\infty(J)} \\ &\lesssim \|R'(T) - \bar{R}'(\bar{T})\|_{L^\infty(J)} + \|\Theta - \bar{\Theta}\|_{L^\infty(J)}. \end{aligned} \quad (4.7)$$

Therefore, the lemma will follow if we establish the estimates:

$$\|R'(T) - \bar{R}'(\bar{T})\|_{L^\infty(J)} \lesssim |J|^{-1} [\|y - \bar{y}\|_{L^\infty(J)} + 2^{-rn}] \quad (4.8)$$

and

$$\|\Theta - \bar{\Theta}\|_{L^\infty(J)} \lesssim |J|^{-1} [\|y - \bar{y}\|_{L^\infty(J)} + 2^{-(r-1)n}]. \quad (4.9)$$

To this end, we need the following estimates:

$$\begin{aligned} \text{(i)} \quad & \|Q(y) - \bar{Q}(\bar{y})\|_{L^\infty(J)} \lesssim \|y - \bar{y}\|_{L^\infty(J)} + 2^{-rn}, \\ \text{(ii)} \quad & \|Q'(y) - \bar{Q}'(\bar{y})\|_{L^\infty(J)} \lesssim \|y - \bar{y}\|_{L^\infty(J)} + 2^{-(r-1)n}, \\ \text{(iii)} \quad & \|T - \bar{T}\|_{L^\infty(J)} \lesssim \|y - \bar{y}\|_{L^\infty(J)} + 2^{-rn}. \end{aligned} \quad (4.10)$$

**Proof of (4.10, i):** Let us denote  $Q_\varepsilon := s^{-1}tg'_n(c \cdot)$ . Then

$$\|Q(y) - \bar{Q}(\bar{y})\|_{L^\infty(J)} \leq \|Q(y) - Q_\varepsilon(\bar{y})\|_{L^\infty(J)} + \|Q_\varepsilon(\bar{y}) - \bar{Q}(\bar{y})\|_{L^\infty(J)}.$$

It follows from (3.4) that

$$\|Q(y) - Q_\varepsilon(\bar{y})\|_{L^\infty(J)} \leq \frac{2\bar{m}}{m} \|y - \bar{y}\|_{L^\infty(J)},$$

and since  $\bar{Q}$  coincides with  $s^{-1}tg'_{n-1}(c \cdot)$  on  $\bar{y}(J)$ , we infer from (3.3) that

$$\|Q_\varepsilon(\bar{y}) - \bar{Q}(\bar{y})\|_{L^\infty(J)} \lesssim 2^{-nr}.$$

**Proof of (4.10, ii):** The same argument can be applied here, since (3.3) implies in particular that  $\|g_n^{(3)}\|_{L^\infty(\Omega)}$  is bounded independantly of  $n$  as long as  $r \geq 2$ .

**Proof of (4.10, iii):** By (4.10, i), we have

$$\begin{aligned} \|T - \bar{T}\|_{L^\infty(J)} &\leq \|y - \bar{y}\|_{L^\infty(J)} + \|Q(y) - \bar{Q}(\bar{y})\|_{L^\infty(J)} \\ &\lesssim 2^{-nr} + \|y - \bar{y}\|_{L^\infty(J)}. \end{aligned}$$

**Proof of (4.8):** Assume first that  $T(J) \cap \bar{T}(J) = \emptyset$ , and without loss of generality, that  $a := \sup(T(J)) < \inf(\bar{T}(J))$ . We extend  $R$  by setting  $R_\varepsilon(x) = R(a) + (x - a)R'(a)$  for  $x \geq a$ . Then

$$\|R'(T) - \bar{R}'(\bar{T})\|_{L^\infty(J)} \leq \|R'(T) - R'_\varepsilon(\bar{T})\|_{L^\infty(J)} + \|R'_\varepsilon(\bar{T}) - \bar{R}'(\bar{T})\|_{L^\infty(J)}.$$

Since  $R'_\varepsilon(\bar{T})$  is a constant over  $J$ , we have

$$\begin{aligned} \|R'(T) - R'_\varepsilon(\bar{T})\|_{L^\infty(J)} &\leq \|R''\|_{L^\infty(T(J))} |T(J)| \\ &\lesssim \|R'\|_{L^\infty(T(J))} \\ &\lesssim 1 \\ &\lesssim |J|^{-1} \|T - \bar{T}\|_{L^\infty(J)}. \end{aligned} \tag{4.11}$$

Here the second inequality is **(P<sub>2</sub>)**, the third inequality is **(A<sub>2</sub>)**, and for the latter inequality, we note that since  $T(J)$  and  $\bar{T}(J)$  are disjoint, then using (4.3),

$$|J| \sim \min(|T(J)|, |\bar{T}(J)|) \leq \|T - \bar{T}\|_{L^\infty(J)}.$$

On the other hand,  $R_e - \bar{R}$  is a polynomial over  $\bar{T}(J)$  and hence we can apply again **(P<sub>2</sub>)** and (4.3) to obtain

$$\begin{aligned} \|R'_e - \bar{R}'\|_{L^\infty(\bar{T}(J))} &\lesssim |J|^{-1} \|R_e - \bar{R}\|_{L^\infty(\bar{T}(J))} \\ &\lesssim |J|^{-1} [\|R_e(\bar{T}) - R(T)\|_{L^\infty(J)} + \|R(T) - \bar{R}(\bar{T})\|_{L^\infty(J)}] \\ &\lesssim |J|^{-1} [\|T - \bar{T}\|_{L^\infty(J)} + \|y - \bar{y}\|_{L^\infty(J)}], \end{aligned}$$

where we used **(A<sub>2</sub>)** and (3.7) for the latter estimate. Together with (4.11) and (4.10, iii), this proves (4.8) in the case where  $T(J)$  and  $\bar{T}(J)$  are disjoint.

Let  $T(J) \cap \bar{T}(J) \neq \emptyset$  and set  $K := T(J) \cup \bar{T}(J)$ . By (4.3),  $K$  is an interval of length  $\mathcal{O}(|J|)$ . Applying **(P<sub>1</sub>)** and **(P<sub>2</sub>)**, we obtain

$$\|R'\|_{L^\infty(K)} \lesssim \|R'\|_{L^\infty(T(J))} \lesssim 1$$

and

$$\|R''\|_{L^\infty(K)} \lesssim |J|^{-1}.$$

We have then

$$\|R'(T) - R'(\bar{T})\|_{L^\infty(J)} \lesssim |J|^{-1} \|T - \bar{T}\|_{L^\infty(J)}$$

and also

$$\begin{aligned} \|R' - \bar{R}'\|_{L^\infty(\bar{T}(J))} &\lesssim |J|^{-1} \|R - \bar{R}\|_{L^\infty(\bar{T}(J))} \\ &\lesssim |J|^{-1} [\|R(T) - R(\bar{T})\|_{L^\infty(J)} + \|R(\bar{T}) - \bar{R}(\bar{T})\|_{L^\infty(J)}] \\ &\lesssim |J|^{-1} [\|R'\|_{L^\infty(K)} \|T - \bar{T}\|_{L^\infty(J)} + \|y - \bar{y}\|_{L^\infty(J)}] \\ &\lesssim |J|^{-1} [\|T - \bar{T}\|_{L^\infty(J)} + \|y - \bar{y}\|_{L^\infty(J)}]. \end{aligned}$$

Consequently,

$$\begin{aligned} \|R'(T) - \bar{R}'(\bar{T})\|_{L^\infty(J)} &\leq \|R'(T) - R'(\bar{T})\|_{L^\infty(J)} + \|R' - \bar{R}'\|_{L^\infty(\bar{T}(J))} \\ &\lesssim |J|^{-1} [\|T - \bar{T}\|_{L^\infty(J)} + \|y - \bar{y}\|_{L^\infty(J)}]. \end{aligned}$$

In view of (4.10, iii), this completes the proof of (4.8).

**Proof of (4.9):** Observe that **(A<sub>2</sub>)** guarantees the boundedness of  $R'$  on  $T(J)$ , and since  $R$  is also obviously bounded on  $T(J)$ , we can apply **(P<sub>2</sub>)** to obtain

$$\|R'\|_{L^\infty(T(J))} \lesssim |J|^{-1}.$$

Then using the definition of  $\Theta$ , we have

$$\begin{aligned}
\|\Theta - \bar{\Theta}\|_{L^\infty(J)} &= \|R'(T)(1 - Q'(y)) - \bar{R}'(\bar{T})(1 - \bar{Q}'(\bar{y}))\|_{L^\infty(J)} \\
&\leq \|R'(T) - \bar{R}'(\bar{T})\|_{L^\infty(J)} \\
&\quad + \|R'(T)\|_{L^\infty(J)} \|Q'(y) - \bar{Q}'(\bar{y})\|_{L^\infty(J)} \\
&\quad + \|\bar{Q}'(\bar{y})\|_{L^\infty(J)} \|R'(T) - \bar{R}'(\bar{T})\|_{L^\infty(J)} \\
&\lesssim |J|^{-1} [\|y - \bar{y}\|_{L^\infty(J)} + 2^{-(r-1)n}],
\end{aligned}$$

where the latter inequality follows from (4.8) and (4.10, ii). This completes the proof of Lemma 4.1.  $\square$

## 4.2 Proof of Lemma 3.2

For simplicity, we denote  $A' := A' \cdot \mathbb{1}_J$  and proceed to estimating  $\|A'\|_{B_{q,q}^{\alpha-1}}$  following the approach of DeVore and Lucier [4]. Recall first the following inverse estimate (Lemma 4.3 in [4]).

**Lemma 4.2.** *Let  $v$  be twice continuously differentiable on an open interval  $I$  and assume that  $v$ ,  $v'$  and  $v''$  each have one sign on  $I$ . If numbers  $p$  and  $q$  are given such that  $0 < p \leq 1$  and  $\frac{1}{p} - \frac{1}{q} > 1$ , then there exists a constant  $C$  such that whenever  $v \in L^q(I)$  then  $v' \in L^p(I)$  and*

$$\|v'\|_{L^p(I)} \leq C |I|^{\frac{1}{p} - \frac{1}{q} - 1} \|v\|_{L^q(I)}. \quad (4.12)$$

According to the definition of the Besov norm in (1.4)-(1.5), we have to estimate  $\omega_k(A', t)_q := \sup_{|h| \leq t} \|\Delta_h^k A'\|_{L^q(\mathbb{R})}$  for  $t > 0$ . Then because of the symmetry, it suffices to bound  $\|\Delta_h^k A'\|_{L^q}$  only for  $0 < h \leq t$ . For a fixed  $h > 0$ , we introduce the following sets:

$$\Gamma := \{x \in \mathbb{R} : [x, x + kh] \subset J\}, \quad \Gamma' := \{x \in \mathbb{R} \setminus \Gamma : [x, x + kh] \cap J \neq \emptyset\},$$

and

$$\Gamma'' := \mathbb{R} \setminus (\Gamma \cup \Gamma') = \{x \in \mathbb{R} : [x, x + kh] \cap J = \emptyset\}.$$

If  $x \in \Gamma''$ , then  $\Delta_h^k A'(x) = 0$  and hence

$$\int_{\Gamma''} |\Delta_h^k A'(x)|^q dx = 0. \quad (4.13)$$

If  $x \in \Gamma'$ , then using that  $|\Delta_h^k A'(x)| \leq 2^k(|A'(x)| + \cdots + |A'(x + kh)|)$ , we have

$$\int_{\Gamma'} |\Delta_h^k A'(x)|^q dx \leq |\Gamma'| \|\Delta_h^k A'\|_{L^\infty(J)}^q \lesssim |\Gamma'| \|A'\|_{L^\infty(J)}^q.$$

Now, Lemma 4.2 and the obvious estimate  $|\Gamma'| \leq \min(h, |J|)$  yield

$$\int_{\Gamma'} |\Delta_h^k A'(x)|^q dx \lesssim \min(h, |J|) |J|^{-q} (\|A\|_{L^\infty(J)} + 2^{-(r-1)n})^q. \quad (4.14)$$

Finally, let  $x \in \Gamma$  and  $0 < h \leq |J|/k$ . Notice that  $\Gamma = \emptyset$  if  $h > |J|/k$ . We shall employ the well-known identity:  $\Delta_h^k A'(x) = h^k A^{(k+1)}(\xi)$  for some  $\xi \in [x, x + kh]$ . From this and the monotonicity of  $A^{(k+1)}$ , we have

$$A^{(k+1)}(\xi) = h^k \max\{A^{(k+1)}(x), A^{(k+1)}(x + kh)\}.$$

Without loss of generality, we can assume that  $A^{(k+1)}$  is decreasing. Then

$$\Delta_h^k A'(x) \leq h^k A^{(k+1)}(x), \quad x \in \Gamma. \quad (4.15)$$

The following embedding is well-known: If  $1 < \beta_1 < \beta_2$ ,  $q_j = 1/\beta_j$ , and  $f \in B_{q_2, q_2}^{\beta_2-1}$ , then  $f \in B_{q_1, q_1}^{\beta_1-1}$  and  $\|f\|_{B_{q_1, q_1}^{\beta_1-1}} \lesssim \|f\|_{B_{q_2, q_2}^{\beta_2-1}}$ . Therefore, we may assume that  $k < \alpha < k+1$ .

Set  $q_0 := q = 1/\alpha$ ,  $\varepsilon := \frac{1}{2}(\frac{\alpha}{k} - 1) > 0$ , and define  $q_1, q_2, \dots, q_k$  recursively by the identity  $\frac{1}{q_j} := \frac{1}{q_{j-1}} - (1 + \varepsilon)$ ,  $j = 1, \dots, k$ . Evidently,  $\frac{1}{q_j} := \alpha - j(1 + \varepsilon)$  and hence  $\frac{1}{q_k} := \alpha - k(1 + \varepsilon) = \frac{1}{2}(\alpha - k) > 0$ . Therefore,  $0 < q_0 < q_1 < \cdots < q_{k-1} < 1$  and  $q_k > 1$ . Now, applying repeatedly Lemma 4.2, we obtain

$$\begin{aligned} \|A^{(k+1)}\|_{L^q(J)} &\lesssim |J|^\varepsilon \|A^{(k)}\|_{L^{q_1}(J)} \lesssim \cdots \lesssim |J|^{k\varepsilon} \|A'\|_{L^{q_k}(J)} \\ &\lesssim |J|^{k\varepsilon + 1/q_k} \|A'\|_{L^\infty(J)} = c |J|^{1/q - \alpha} \|A'\|_{L^\infty(J)}. \end{aligned} \quad (4.16)$$

Using (4.15), (4.16), and Lemma 4.1, we get

$$\int_{\Gamma} |\Delta_h^k A'(x)|^q dx \lesssim h^{kq} |J|^{1-q-kq} (\|A\|_{L^\infty(J)} + 2^{-(r-1)n})^q. \quad (4.17)$$

Combining (4.13), (4.14), and (4.17), we arrive at

$$\begin{aligned} \omega_k(A', t)_q^q &= \sup_{0 < h \leq t} \int_{\mathbb{R}} |\Delta_h^k A'(x)|^q dx \\ &\lesssim [\min(t, |J|) + t^{kq} |J|^{1-kq} \mathbf{1}(t)] |J|^{-q} (\|A\|_{L^\infty(J)} + 2^{-(r-1)n})^q, \end{aligned}$$



where  $\mathbb{1} := \mathbb{1}_{[0, |J|/k]}$ . Therefore,

$$\begin{aligned} \|A'\|_{B_{q,q}^{\alpha-1}}^q &= \int_0^\infty t^{-(\alpha-1)q-1} \omega_k(A', t)^q dt \\ &\lesssim [ |J|^{-q} \int_0^{|J|} t^{q-1} dt + |J|^{1-q} \int_{|J|}^\infty t^{q-2} dt \\ &\quad + |J|^{1-q-kq} \int_0^{|J|/k} t^{q+kq-2} dt ] (\|A\|_{L^\infty(J)} + 2^{-(r-1)n})^q \\ &\lesssim (\|A\|_{L^\infty(J)} + 2^{-(r-1)n})^q, \end{aligned}$$

where we used that  $0 < q < 1$  and  $kq + q - 2 = (k+1)/\alpha - 2 > -1$ . The proof of Lemma 3.2 is complete.  $\square$

## References

- [1] M. Campos-Pinto, A. Cohen, W. Dahmen and R. A. DeVore, *On the stability of nonlinear conservation laws in the Hausdorff metric*, preprint Laboratoire J.-L. Lions, Université Pierre et Marie Curie, Paris, 2003.
- [2] R. A. DeVore, *Nonlinear approximation*, Acta numerica, No. 7, Cambridge Univ. Press, Cambridge, 1998, 51-150.
- [3] R. A. DeVore and B. J. Lucier, *High order regularity for solutions of the inviscid Burgers equation*, in “Nonlinear hyperbolic problems (Bordeaux, 1988)”, Lecture Notes in Math., 1402, Springer, Berlin, 1989, 147-154.
- [4] R. A. DeVore and B. J. Lucier, *High order regularity for conservation laws*, Indiana Math. J. 39, No. 2, 1990, 413-430.
- [5] R. A. DeVore and V. A. Popov, *Interpolation spaces and non-linear approximation*, in “Function spaces and applications”, Lecture Notes in Math., 1302, Springer, Berlin, 1988, 191-205.
- [6] E. Godlewski and P.-A. Raviart, *Numerical approximation of hyperbolic systems of conservation laws*, Applied Mathematical Sciences, 118. Springer-Verlag, New York, 1996.
- [7] D. Kröner, *Numerical schemes for conservation laws*, Advances in numerical mathematics, Wiley-Teubner, 1997.
- [8] S.N. Kruzkov, *First order quasilinear equations in several independent variables*, Math. USSR Sb. 10, 1970, 217-243.

- [9] P. D. Lax, *Hyperbolic systems of conservation laws and the mathematical theory of shock waves*, in “Regional Conference Series in Applied Mathematics”, No. 11. SIAM, Philadelphia, 1973.
- [10] R. J. LeVeque, *Numerical methods for conservation laws*, Lectures in Mathematics ETH Zurich. Birkhauser Verlag, Basel, 1990.
- [11] B. J. Lucier, *A moving mesh numerical method for hyperbolic conservation laws*, Math. Comp. 46, 1986, No. 173, 59-69.
- [12] P. Petrushev, *Direct and converse theorems for spline and rational approximation and Besov spaces*, in “Function spaces and applications”, Lecture Notes in Math., 1302, Springer, Berlin, 1988, 363-377.
- [13] Bl. Sendov, *Hausdorff Approximations*, Kluver Acad. Pub., 1990.